

On the non-existence of certain hyperovals in dual André planes of order 2^{2h}

A. Aguglia and L. Giuzzi*

Abstract

No regular hyperoval of the Desarguesian affine plane $AG(2, 2^{2h})$, with $h > 1$, is inherited by a dual André plane of order 2^{2h} with dimension 2 over its centre.

1 Introduction

The general question on existence of ovals in finite non-Desarguesian planes is still open and appears to be difficult. It has been shown by computer search that there exist some planes of order 16 without ovals; see [11]. On the other hand, ovals have been constructed in several finite planes; one of the most fruitful approaches in this search has been that of inherited oval, due to Korchmáros [5, 6].

Korchmáros' idea relies on the fact that any two planes π_1 and π_2 of the same order have the same number of points and lines; thus their point sets, as well as some lines, may be identified. If Ω is an oval of π_1 , it might happen that Ω , regarded as a point set, turns also out to be an oval of π_2 , although π_1 and π_2 differ in some (in general several) point-line incidences; in this case Ω is called an *inherited oval* of π_2 from π_1 ; see also [2, Page 728].

In practice, it is usually convenient to take π_1 to be the Desarguesian affine plane $AG(2, q)$ of order a prime power q . The case in which π_2 is the Hall plane $H(q^2)$ of order q^2 was investigated in [5], and inherited ovals were found. For q odd, this also proves the existence of inherited ovals in the dual plane of $H(q^2)$, which is a Moulton plane $M(q^2)$ of the same order.

Moulton planes have been originally introduced in [10], by altering some of the lines of a Desarguesian plane constructed over the reals, while keeping the original point set fixed. In particular, each line of the Moulton plane turns out to be either a line of the original plane or the union of two half-lines of different slope with one point in common.

*Research supported by the Italian Ministry MIUR, Strutture geometriche, combinatoria e loro applicazioni.

This construction, when considering planes of finite order q^2 , may be carried out as follows. Let $\|\cdot\|$ denote the norm function

$$\|\cdot\| : \begin{cases} \text{GF}(q^2) \rightarrow \text{GF}(q) \\ x \mapsto x^{q+1} \end{cases}$$

Take a proper subset U of $\text{GF}(q)^*$ and consider the following operation defined over the set $\text{GF}(q^2)$

$$a \odot b = \begin{cases} ab & \text{if } \|b\| \notin U \\ a^q b & \text{if } \|b\| \in U. \end{cases}$$

The set $(\text{GF}(q^2), +, \odot)$ is a pre-quasifield which is a quasifield for $U \neq \{1\}$. Every pre-quasifield coordinatizes a translation plane; see [4, Section 5.6]. In our case this translation plane is an affine André plane $A(q^2)$ of order q^2 with dimension 2 over its centre. In the case in which U consists of a single element of $\text{GF}(q^2)$ the translation plane is the affine Hall plane of order q^2 and its dual plane is the affine Moulton plane of order q^2 . For details on these planes see [3, 8].

Write $\mathcal{M}_U(q^2) = (\mathcal{P}, \mathcal{L})$ for the incidence structure whose the point-set $\mathcal{P}$ is the same as that of $AG(2, q^2)$, and whose lines in $\mathcal{L}$ are either of the form

$$[c] = \{P(x, y) : x = c, y \in \text{GF}(q^2)\}$$

or

$$[m, n] = \{P(x, y) : y = m \odot x + n\}.$$

The affine plane $\mathcal{M}_U(q^2)$ is the dual of an affine André plane $A(q^2)$ of order q^2 . Completing $\mathcal{M}_U(q^2)$ with its points at infinity in the usual way gives a projective plane $\overline{\mathcal{M}_U(q^2)}$ called *the projective closure of $\mathcal{M}_U(q^2)$* .

Write $\Phi = \{P(x, y) : \|x\| \notin U\}$ and $\Psi = \{P(x, y) : \|x\| \in U\}$. Clearly, $\mathcal{P} = \Phi \cup \Psi$.

If an arc $\mathcal{A}$ of $PG(2, q^2)$ is in turn an arc in $\overline{\mathcal{M}_U(q^2)}$ then, $\mathcal{A}$ is an *inherited arc* of $\overline{\mathcal{M}_U(q^2)}$.

Any hyperoval of the Desarguesian projective plane $PG(2, q^2)$ obtained from a conic by adding its nucleus is called *regular*. Let consider a regular hyperoval Ω in $AG(2, q^2)$. If $\Omega \subseteq \Phi$, that is for each point $P(x, y) \in \Omega$ the norm of x is an element of $\text{GF}(q) \setminus U$, then Ω is clearly an inherited hyperoval of $\mathcal{M}_U(q^2)$.

In [1, Theorem 1.1], it is proven that for $q > 5$ an odd prime power, any arc of the Moulton plane $\mathcal{M}_t(q^2)$ with $t \in \text{GF}(q)$, obtained as $\mathcal{C}^* = \mathcal{C} \cap \Phi$, where $\mathcal{C}$ is an ellipse in $AG(2, q^2)$ is complete.

In this paper the case where q is even and $|U| < \frac{q}{4} - 1$ is addressed. We prove the following.

Theorem 1. *Suppose Ω to be a regular hyperoval of $AG(2, 2^{2h})$, with $h > 1$. Then, $\Omega^* = \Omega \cap \Phi$ is a complete arc in the projective closure of $\mathcal{M}_U(2^{2h})$.*

Theorem 2. *No regular hyperoval Ω of $AG(2, 2^{2h})$, with $h > 1$, is an inherited hyperoval of $\mathcal{M}_U(2^{2h})$.*

We shall also see that any oval arising from a regular hyperoval of $AG(2, 2^{2h})$ by deleting a point cannot be inherited by $\mathcal{M}_U(2^{2h})$. The hypothesis on Ω being a regular hyperoval cannot be dropped; see [11] for examples of hyperovals in the Moulton plane of order 16.

2 Proof of Theorem 1

We begin by showing the following lemma, which is a slight generalisation of Lemma 2.1. in [1].

Lemma 3. *Let q be any prime power. A pencil of affine lines $\mathcal{L}(P)$ of $\mathcal{M}_U(q^2)$ with centre $P(x_0, y_0)$, either consists of lines of a Baer subplane $\mathcal{B}$ of $PG(2, q^2)$, or is a pencil in $AG(2, q^2)$ with the same center according to $\|x_0\| \in U$ or not. In particular, in the former case, the $q^2 + 1$ lines in $\mathcal{L}(P)$ plus the q vertical lines $X = c$ with $\|c\| = x_0^{q+1}$ and $c \neq x_0$ are the lines of $\mathcal{B}$.*

Proof. The pencil $\mathcal{L}(P)$ consists of the lines

$$r_m : y = m \odot x - m \odot x_0 + y_0,$$

with $m \in GF(q^2)$, plus the vertical line $\ell : x = x_0$. First suppose $\|x_0\| \in U$. In this case $m \odot x_0 = m^q x_0$ and the line r_m of $\mathcal{L}(P)$ corresponds to the point $(m, m^q x_0 - y_0)$ in the dual of $\mathcal{M}_U(q^2)$, which is an André plane.

As m varies over $GF(q^2)$ we get q^2 affine points of the Baer subplane $\mathcal{B}'$ in $PG(2, q^2)$ represented by $y = x^q x_0 - y_0$. The points at infinity of $\mathcal{B}'$ are those points (c) such that $c^{q+1} = \|x_0\|$. As the dual of a Baer subplane is a Baer subplane, it follows that the lines in $\mathcal{L}(P)$ are the lines of a Baer subplane $\mathcal{B}$ in $PG(2, q^2)$. More precisely, the lines in $\mathcal{L}(P)$ plus the q vertical lines $x = c$, $\|c\| = \|x_0\|$, with $c \neq x_0$, are the lines of $\mathcal{B}$.

In the case in which $\|x_0\| \notin U$ the line $r_m : y = m \odot x - mx_0 + y_0$ in $\mathcal{L}(P)$ corresponds to the point

$$(m, mx_0 - y_0)$$

in the dual of $\mathcal{M}_U(q^2)$. As m varies over $GF(q^2)$ we get q^2 affine points in $AG(2, q^2)$ on the line $y = x_0 x - y_0$. The dual of each such a point is a line of $AG(2, q^2)$ through the point $P(x_0, y_0)$. $\square$

Let Ω denote a regular hyperoval in $AG(2, q^2)$, $q = 2^h$, $h > 1$. It will be shown that for any point $P(x_0, y_0)$ with $\|x_0\| \in U$ there is at least a 2-secant to $\Omega^* = \Omega \cap \Phi$ in $\mathcal{M}_U(q^2)$ through P .

Assume $\mathcal{B}$ to be the Baer subplane in $PG(2, q^2)$ containing the lines of $\mathcal{L}(P)$ and the q vertical lines $X = c$ with $\|c\| = x_0^{q+1}$, $c \neq x_0$. Write Δ for the set of all points of Ω not covered by a vertical line of $\mathcal{B}$ and also let $n = |\Delta|$ and $m = q^2 + 2 - n$. The vertical lines of $\mathcal{B}$ cover at most $2(q+1)$ points of Ω ; thus, $q^2 - 2q \leq n \leq q^2 + 2$. We shall show that there is at least a line in $\mathcal{B}$ meeting Δ in two points.

Let $T \in \Delta$; since $T \notin \mathcal{B}$, there is a unique line ℓ_T of $\mathcal{B}$ through T . Every point $Q \in \Omega \setminus \Delta$ lies on at most $q+1-(m-1) = q-m+2$ lines ℓ_T with $T \in \Delta$. Suppose by contradiction that for every $T \in \Delta$,

$$\ell_T \cap \Omega = \{T, Q\}, \text{ with } Q \in \Omega \setminus \Delta.$$

The total number of lines ℓ_T obtained as Q varies in $\Omega \setminus \Delta$ does not exceed $m(q-m+2)$. So,

$$n = q^2 - m + 2 \leq m(q-m+2).$$

As m is a non-negative integer, this is possible only for $q = 2$.

Since ℓ_T is not a vertical line, it turns out to be a chord of Ω^* in $\mathcal{M}_U(q^2)$ passing through $P(x_0, y_0)$. This implies that no point $P(x_0, y_0) \in \Psi$ may be aggregated to Ω^* in order to obtain an arc.

This holds true in the case $P(x_0, y_0) \in \Phi$. In $AG(2, q^2)$ there pass $(q^2+2)/2$ secants to Ω through a point $P(x_0, y_0) \notin \Omega$ and, hence, $N = (q^2+2)/2 - s$ secants to Ω^* , where $s \leq 2(q+1)|U|$. So by the hypothesis $|U| < q/4 - 1$, we obtain $N > 0$; this implies that no point $P(x_0, y_0) \in \Phi$ may be aggregated to Ω^* in order to obtain a larger arc. The same argument works also when P is assumed to be a point at infinity. Theorem 1 is thus proved.

3 Proof of Theorem 2

We shall use the notion of *conic blocking set*; see [7]. A conic blocking set $\mathcal{B}$ is a set of lines in a Desarguesian projective plane met by all conics; a conic blocking set $\mathcal{B}$ is *irreducible* if for any line of $\mathcal{B}$ there is a conic intersecting $\mathcal{B}$ in just that line.

Lemma 4 (Theorem 4.4, [7]). *The line-set*

$$\mathcal{B} = \{y = mx : m \in \text{GF}(q)\} \cup \{x = 0\}$$

is an irreducible conic blocking set in $\text{PG}(2, q^2)$, where $q = 2^h$, $h > 1$.

Lemma 5. *Let Ω be a regular hyperoval of $\text{PG}(2, q^2)$, with $q = 2^h$, $h > 1$. Then, there are at least two points $P(x, y)$ in Ω such that $\|x\| \in U$.*

Proof. To prove the lemma we show that the set $\Psi' = \Psi \cup Y_\infty$, is a conic blocking set. We observe that the conic blocking set of Lemma 4 is actually a degenerate Hermitian curve of $\text{PG}(2, q^2)$ with equation $x^q y - x y^q = 0$. Since all degenerate Hermitian curves are projectively equivalent, this implies that any such a curve is a conic blocking set. On the other hand, Ψ' may be regarded as the union of degenerate Hermitian curves of equation $x^{q+1} = c z^{q+1}$, as c varies in U . Thus, Ψ' is also a conic blocking set. Suppose now $\Omega = \mathcal{C} \cup N$, where $\mathcal{C}$ is a conic of nucleus N . Take $P \in \Psi' \cap \mathcal{C}$. If $P = Y_\infty$ then at most one of the vertical lines $X = c$, with $\|c\| \in U$, is tangent to $\mathcal{C}$; hence there are at least q points $P'(x, y) \in \Omega$ with $\|x\| \in U$ and thus $|\Psi \cap \Omega| \geq q$.

Next, assume that $P = P(x, y) \in \Psi$. If the line $[x]$ is secant to $\mathcal{C}$ the assertion immediately follows. If the line $[x]$ is tangent to $\mathcal{C}$ then the nucleus N lies on $[x]$. Now, either N is an affine point in Ψ or $N = Y_\infty$. In the former case we have $|\Psi \cap \Omega| \geq 2$; in the latter, the lines $X = c$ with $\|c\| \in U$ are all tangent to $\mathcal{C}$; hence, there are at least other $q + 1$ points $P'(x, y) \in \Omega$ such that $\|x\| \in U$. $\square$

Now, let Ω be a regular hyperoval in $AG(2, q^2)$, with $q = 2^h$ and $h > 1$. From Lemma 5 we deduce that $|\Omega^* \cap \Phi| \leq q^2$; furthermore, Theorem 2 guaranties that Ω^* is a complete arc in the projective closure of $\mathcal{M}_U(q^2)$, whence Theorem 2 follows.

Remark 1. The largest arc of $\mathcal{M}_U(q^2)$ contained in a regular hyperoval of $AG(2, q^2)$, with $q = 2^h$, has at most q^2 points; in particular any oval which arises from a hyperoval of $AG(2, q^2)$ by deleting a point cannot be an oval of $\mathcal{M}_U(q^2)$. For an actual example of a q^2 -arc of $\mathcal{M}_U(q^2)$ coming from a regular hyperoval of $AG(2, q^2)$ see [9]. This also shows that the result of [5] cannot be extended to even q .

References

- [1] V. Abatangelo, B. Larato, *Canonically Inherited Arcs in Moulton Planes of Odd Order*, to appear on Innovations in Incidence Geometry.
- [2] C.J. Colbourn, J.H. Diniz, *Handbook of Combinatorial Designs*, Chapman & Hall (2007).
- [3] P. Dembowski, *Finite Geometries*, Springer (1968).
- [4] N.L. Johnson, V. Jha, M. Biliotti, *Handbook of finite translation planes*. Pure and Applied Mathematics (Boca Raton), **289** Chapman & Hall/CRC, Boca Raton, FL, (2007).
- [5] G. Korchmáros, *Ovali nei piani di Moulton d'ordine dispari*, Colloquio Internazionale sulle Teorie Combinatorie (Rome, 1973), Tomo II, 395–398. Atti dei Convegni Lincei, No. **17**, Accad. Naz. Lincei, Rome, (1976).
- [6] G. Korchmáros, *Ovali nei piani di Hall di ordine dispari*, Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur. (8) **56** (1974), no. 3, 315–317.
- [7] L. D. Holder, *Conic blocking sets in Desarguesian projective planes*, J. Geom. **80** (2004), no. 1-2, 95–105.
- [8] D. Hughes, F. Piper, *Projective planes*, Springer (1973).
- [9] G. Menichetti, *k-archi completi in piani di Moulton d'ordine q^m* , Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur. (8) **60** (1976), no. 6, 775–781.

- [10] F.R. Moulton, *A simple non-desarguesian plane geometry*, Trans. Am. Math. Soc. **3** (1902), 192–195.
- [11] T. Penttila, G.F. Royle, M.K. Simpson, *Hyperovals in the known projective planes of order 16*, J. Combin. Des. **4** (1996), no. 1, 59–65.

ANGELA AGUGLIA
Dipartimento di Matematica
Politecnico di Bari
Via Orabona 4
I-70125 Bari
Italy
e-mail: a.aguglia@poliba.it

LUCA GIUZZI
Dipartimento di Matematica
Università degli Studi di Brescia
Facoltà di Ingegneria
Via Valotti 9
I-25133 Brescia
Italy
e-mail: giuzzi@ing.unibs.it